

# Immune Globulin Intravenous (Human) Panglobulin®

## Lyophilized Preparation

<b>SAMPLE INSTRUCTIONS ONLY</b> Directions for use are provided with the product and may be frequently revised. This insert is for educational purposes only. When performing laboratory techniques or using the product in therapeutic situations, use only package insert supplied with the product.
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

## Rx only Description

Immune Globulin Intravenous (Human) (IGIV), Panglobulin®, is a sterile, highly purified polyvalent antibody product containing in concentrated form all the IgG antibodies which regularly occur in the donor population (1). This immunoglobulin preparation is produced by cold alcohol fractionation from the plasma of volunteer US donors. Part of the fractionation may be performed by another US-licensed manufacturer. The fractionation process by which Panglobulin® is prepared from plasma includes several filtration steps which are carried out in the presence of filter aids; some of these filtration steps are used for the separation of a cold ethanol precipitate. Four of these steps were validated for virus elimination. The cumulative LRFs (log<sub>10</sub> of reduction factors) were 15.5 for HIV (human immunodeficiency virus), 16.0 for PRV (pseudorabies virus), 9.3 for SFV (Semliki Forest virus), 12.4 for Sindbis virus, and 14.1 for BEV (bovine enterovirus).

Panglobulin® is made suitable for intravenous use by treatment at acid pH in the presence of trace amounts of pepsin (2,3). Treatment with pepsin at pH4 rapidly inactivates enveloped viruses. LRFs were ≥ 6.1 for HIV, ≥ 5.3 for PRV, ≥ 4.4 for BVDV (bovine viral diarrhea virus), and ≥ 6.8 for SFV. PRV and the two model viruses for HCV (hepatitis C virus), BVDV and SFV, were all inactivated within 1/10, and HIV within 1/2 of the total incubation time used during production of Panglobulin®. Overall viral clearance by either elimination and/or inactivation during the manufacturing process has been documented to be ≥21 for HIV, ≥19 for PRV, ≥15 for SFV, and ≥14 for BEV (expressed as LRF). The preparation contains at least 96% of IgG and after reconstitution with a neutral unbuffered diluent has a pH of 6.6 ± 0.2. Most of the immunoglobulins are monomeric (7 S) IgG; the remainder consists of dimeric IgG and a small amount of polymeric IgG, traces of IgA and IgM and immunoglobulin fragments (4). The distribution of the IgG subclasses corresponds to that of normal serum (5,6,7,8). Final container lyophilized units are prepared so as to contain 1, 3, 6, or 12 g protein with 1.67 g sucrose and less than 20 mg NaCl per gram of protein. The lyophilized preparation is devoid of any preservatives and may be reconstituted with sterile water, 5% dextrose or 0.9% saline to a solution with protein concentrations ranging from 3% to 12%.

The patient's fluid, electrolyte, caloric requirements and renal function should be considered in selecting an appropriate diluent and concentration.

Table 1

#### Calculated Panglobulin® Osmolality (mOsm/kg)

	<b>Concentration</b>			
<b>Diluent</b>	<b>3%</b>	<b>6%</b>	<b>9%</b>	<b>12%</b>
0.9% NaCl	498	690	882	1074
5% Dextrose	444	636	828	1020
Sterile Water	192	384	576	768

## Clinical Pharmacology

This product contains a broad spectrum of antibody specificities against bacterial, viral, parasitic, and mycoplasma antigens, that are capable of both opsonization and neutralization of microbes and toxins. The 3 week half-life of Immune Globulin Intravenous (Human), Panglobulin®, corresponds to that of Immune Globulin (Human) for intramuscular use, although individual variations in half-life have been observed (9,10). Appropriate doses of Panglobulin® restore abnormally low immunoglobulin G levels to the normal range. One hundred percent of the infused dose is available in the recipient's circulation immediately after infusion. After approximately 6 days, an equilibrium is reached between the intra- and extravascular compartments, with immunoglobulin G being distributed approximately 50% intravascular and 50% extravascular. In comparison, after the intramuscular injection of immune globulin, the IgG requires 2–5 days to reach its maximum concentration in the intravascular compartment. This concentration corresponds to about 40% of the injected dose (10).

While Panglobulin® has been shown to be effective in some cases of Immune Thrombocytopenic Purpura (ITP) (see INDICATIONS AND USAGE), the mechanism of action in ITP has not been fully elucidated.

Toxicity from overdose has not been observed on regimens of 0.4 g/kg body weight each day for 5 days (11,12,13). Sucrose is added to Panglobulin® for reasons of stability and solubility.

Since sucrose is excreted unchanged in the urine when given intravenously, Panglobulin®, may be given to diabetics without compensatory changes in insulin dosage regimen. Please see WARNINGS section.

## Indications and Usage

#### Immunodeficiency

Immune Globulin Intravenous (Human), Panglobulin®, is indicated for the maintenance treatment of patients with primary immunodeficiencies, e.g., in common variable immunodeficiency, severe combined immunodeficiency, and primary immunoglobulin deficiency syndromes such as X-linked agammaglobulinemia (12,14,15,16). Panglobulin® is preferable to intramuscular Immune Globulin (Human) preparations in treating patients who require an immediate and large increase in the intravascular immunoglobulin level (10), in patients with limited muscle mass, and in patients with bleeding tendencies for whom intramuscular injections are contraindicated. The infusions must be repeated at regular intervals. Please see DOSAGE AND ADMINISTRATION section.

#### Immune Thrombocytopenic Purpura (ITP)

#### Acute

A controlled study was performed in children in which Immune Globulin Intravenous (Human), Panglobulin®, was compared with steroids for the treatment of acute (defined as less than 6 months duration) ITP. In this study sequential platelet levels of 30,000, 100,000, and 150,000/μl were all achieved faster with Panglobulin® than with steroids and without any of the side effects associated with steroids (11,17). However, it should be noted that many cases of acute ITP in childhood resolve spontaneously within weeks to months. Panglobulin® has been used with good results in the treatment of acute ITP in adult patients (18,19,20). In a study involving 10 adults with ITP of less than 16 weeks duration, Panglobulin® therapy raised the platelet count to the normal range after a 5 day course. This effect lasted a mean of over 173 days, ranging from 30 to 372 days (21).

#### Chronic

Children and adults with chronic (defined as greater than 6 months duration) ITP have also shown an increase (sometimes temporary) in platelet counts upon administration of Immune Globulin Intravenous (Human), Panglobulin®, (17,21,22,23,24,25). Therefore, in situations that require a rapid rise in platelet count, for example prior to surgery or to control excessive bleeding, use of Panglobulin® should be considered. In children with chronic ITP, Panglobulin® therapy resulted in a mean rise in platelet count of 312,000/μl with a duration of increase ranging from 2 to 6 months (22,25). Panglobulin® therapy may be considered as a means to defer or avoid splenectomy (24,25,26). In adults, Panglobulin® therapy has been shown to be effective in maintaining the platelet count in an acceptable range with or without periodic booster therapy. The mean rise in platelet count was 93,000/μl and the average duration of the increase was 20–24 days (21,22). However, it should be noted that not all patients will respond. Even in those patients who do respond, this treatment should not be considered to be curative.

## Contraindications

As with all blood products containing IgA, Immune Globulin Intravenous (Human), Panglobulin®, is contraindicated in patients with selective IgA deficiency, who possess antibody to IgA. It may also be contraindicated in patients who have had severe systemic reactions to the intravenous or intramuscular administration of human immune globulin.

## Warnings

**WARNING**  
**Immune Globulin Intravenous (Human) (IGIV) products have been reported to be associated with renal dysfunction, acute renal failure, osmotic nephrosis, and death (27,28,29,30,31,32). Patients predisposed to acute renal failure include patients with any degree of pre-existing renal insufficiency, diabetes mellitus, age greater than 65, volume depletion, sepsis, paraproteinemia, or patients receiving known nephrotoxic drugs. Especially in such patients, IGIV products should be administered at the minimum concentration available and the minimum rate of infusion practicable. While these reports of renal dysfunction and acute renal failure have been associated with the use of many of the licensed IGIV products, those containing sucrose as a stabilizer accounted for a disproportionate share of the total number.**

**See PRECAUTIONS and DOSAGE AND ADMINISTRATION sections for important information intended to reduce the risk of acute renal failure.**

**Immune Globulin Intravenous (Human), Panglobulin®, is made from human plasma. Products made from human plasma may contain infectious agents, such as viruses, that can cause disease. The risk that such products will transmit an infectious agent has been reduced by screening plasma donors for prior exposure to certain viruses, by testing for the presence of certain current virus infections, and by inactivating and/or removing certain viruses. The fractionation process by which Panglobulin® is prepared from plasma includes several filtration steps which are carried out in the presence of filter aids; some of these filtration steps are used for the separation of a cold ethanol precipitate. Four of these steps were validated for virus elimination. The cumulative LRFs (log<sub>10</sub> of reduction factors) were 15.5 for HIV (human immunodeficiency virus), 16.0 for PRV (pseudorabies virus), 9.3 for SFV (Semliki Forest virus), 12.4 for Sindbis virus, and 14.1 for BEV (bovine enterovirus).**

**Panglobulin® is made suitable for intravenous use by treatment at acid pH in the presence of trace amounts of pepsin (2,3). Treatment with pepsin at pH4 rapidly inactivates enveloped viruses. LRFs were ≥ 6.1 for HIV, ≥ 5.3 for PRV, ≥ 4.4 for BVDV (bovine viral diarrhea virus), and ≥ 6.8 for SFV. PRV and the two model viruses for HCV (hepatitis C virus), BVDV and SFV, were all inactivated within 1/10, and HIV within 1/2 of the total incubation time used during production of Panglobulin®. Overall viral clearance by either elimination and/or inactivation during the manufacturing process has been documented to be ≥ 21 for HIV, ≥ 19 for PRV, ≥ 15 for SFV, and ≥ 14 for BEV (expressed as LRF). Despite these measures, such products may carry a risk of transmitting infectious agents, e.g., viruses, and theoretically, the Creutzfeldt-Jakob disease (CJD) agent. There is also the possibility that unknown infectious agents may be present in such products. ALL infections thought by a physician possibly to**

**have been transmitted by this product should be reported by the physician or other healthcare provider to American Red Cross, Tel. no. 1-800 293 5023. The physician should discuss the risks and benefits of this product with the patient.**

Patients with agamma- or extreme hypogammaglobulinemia who have never before received immunoglobulin substitution treatment or whose time from last treatment is greater than 8 weeks, may be at risk of developing inflammatory reactions on rapid infusion of Immune Globulin Intravenous (Human), Panglobulin®, (over 20 drops [1 mL] per minute). These reactions are manifested by a rise in temperature, chills, nausea, and vomiting. The patient's vital signs should be monitored continuously. The patient should be carefully observed throughout the infusion, since these reactions on rare occasions may lead to shock. Epinephrine should be available for treatment of an acute anaphylactic reaction.

## Precautions

Please see DOSAGE AND ADMINISTRATION below, for important information on Immune Globulin Intravenous (Human), Panglobulin®, compatibility with other medications or fluids.

Assure that patients are not volume depleted prior to the initiation of the infusion of IGIV.

Periodic monitoring of renal function tests and urine output is particularly important in patients judged to have a potential increased risk for developing acute renal failure. Renal function, including measurement of blood urea nitrogen (BUN)/serum creatinine, should be assessed prior to the initial infusion of Panglobulin® and again at appropriate intervals thereafter. If renal function deteriorates, discontinuation of the product should be considered. For patients judged to be at risk for developing renal dysfunction, it may be prudent to reduce the amount of product infused per unit time by infusing Panglobulin® at a rate less than 2 mg Ig/kg/min.

#### Information for Patients

Patients should be instructed to immediately report symptoms of decreased urine output, sudden weight gain, fluid retention/edema, and/or shortness of breath (which may suggest kidney damage) to their physicians.

#### Pregnancy

Pregnancy Category C: Animal reproduction studies have not been conducted with Immune Globulin Intravenous (Human), Panglobulin®. It is also not known whether Panglobulin® can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Panglobulin® should be given to a pregnant woman only if clearly needed (20).

Intact immune globulins such as those contained in Panglobulin® cross the placenta from maternal circulation increasingly after 30 weeks gestation (33,34). In cases of maternal ITP where Panglobulin® was administered to the mother prior to delivery, the platelet response and clinical effect were similar in the mother and neonate (20,34–43).

#### Pediatric Use

High dose administration of Immune Globulin Intravenous (Human), Panglobulin®, in pediatric patients with acute or chronic Immune Thrombocytopenic Purpura did not reveal any pediatric-specific hazard (11).

Antibodies in Immune Globulin Intravenous (Human) may impair the efficacy of live attenuated viral vaccines such as measles, rubella, and mumps (44,45,46). Immunizing physicians should be informed of recent therapy with Immune Globulin Intravenous (Human) so that appropriate precautions may be taken.

#### Aseptic Meningitis Syndrome

An aseptic meningitis syndrome (AMS) has been reported to occur infrequently in association with Immune Globulin Intravenous (Human) (IGIV) treatment. The syndrome usually begins within several hours to two days following IGIV treatment. It is characterized by symptoms and signs including severe headache, nuchal rigidity, drowsiness, fever, photophobia, painful eye movements, and nausea and vomiting. Cerebrospinal fluid studies are frequently positive with pleocytosis up to several thousand cells per cu.mm., predominantly from the granulocytic series, and elevated protein levels up to several hundred mg/dl. Patients exhibiting such symptoms and signs should receive a thorough neurological examination, including CSF studies, to rule out other causes of meningitis. AMS may occur more frequently in association with high dose (2 g/kg) IGIV treatment. Discontinuation of IGIV treatment has resulted in remission of AMS within several days without sequelae.

## Adverse Reactions

Increases in creatinine and blood urea nitrogen (BUN) have been observed as soon as one to two days following infusion. Progression to oliguria or anuria, requiring dialysis has been observed.

Types of severe renal adverse events that have been seen following IGIV therapy include: acute renal failure, acute tubular necrosis, proximal tubular nephropathy and osmotic nephrosis (27-32,47,57–59). Inflammatory adverse reactions have been described in agammaglobulinemic and hypogammaglobulinemic patients who have never received immunoglobulin substitution therapy before or in patients whose time from last treatment is greater than 8 weeks and whose initial infusion rate exceeds 20 drops (1 ml) per minute. This occurs in approximately 10% of such cases. Such reactions may also be observed in some patients during chronic substitution therapy. These reactions, which generally become apparent only 30 minutes to 1 hour after the beginning of the infusion, are as follows: flushing of the face, feelings of tightness in the chest, chills, fever, dizziness, nausea, diaphoresis, and hypotension. In such cases the infusion should be temporarily stopped until the symptoms have subsided. Immediate anaphylactoid and hypersensitivity reactions due to previous sensitization of the recipient to certain antigens, most commonly IgA, may be observed in exceptional cases, described under CONTRAINDICATIONS (12,13,48). In patients with ITP, who receive higher doses (0.4 g/kg/day or greater), 2.9% of infusions may result in adverse reactions (17). Headache, generally mild, is the most common symptom noted, occurring during or following 2% of infusions.

A few cases of usually mild hemolysis have been reported after infusion of intravenous immunoglobulin products (49,50). These were attributed to transferral of blood group (e.g., anti-D) antibodies (51,52).

## Dosage and Administration

It is generally advisable not to dilute plasma derivatives with other infusable drugs. Immune Globulin Intravenous (Human), Panglobulin<sup>®</sup>, should be given by a separate infusion line. No other medications or fluids should be mixed with the Panglobulin<sup>®</sup> preparation.

Panglobulin<sup>®</sup> should be used with caution in patients with pre-existing renal insufficiency and in patients judged to be at increased risk of developing renal insufficiency (including, but not limited to those with diabetes mellitus, age greater than 65, volume depletion, para-proteinemia, sepsis, and patients receiving known nephrotoxic drugs).

In these cases especially it is important to assure that patients are not volume depleted prior to Panglobulin<sup>®</sup> infusion. No prospective data are presently available to identify a maximum safe dose, concentration, and rate of infusion in patients determined to be at increased risk of acute renal failure. In the absence of prospective data, recommended doses should not be exceeded and the concentration and infusion rate selected should be the minimum practicable. The product should be infused at a rate less than 2 mg Ig/kg/min.

#### Adult and Child Substitution Therapy

The usual dose of Immune Globulin Intravenous (Human), Panglobulin<sup>®</sup> in immunodeficiency syndromes is 0.2 g/kg of body weight administered once a month by intravenous infusion. If the clinical response is inadequate, the dose may be increased to 0.3 g/kg of body weight or the infusion may be repeated more frequently than once a month (12,14,15,16).

The first infusion of Panglobulin<sup>®</sup> in previously untreated agammaglobulinemic or hypogammaglobulinemic patients must be given as a 3% immunoglobulin solution (use the total volume of fluid provided, or see Table 2, to reconstitute the lyophilized product). Start with a flow rate of 10–20 drops (0.5–1.0 mL) per minute. After 15–30 minutes the rate of infusion may be further increased to 30–50 drops (1.5–2.5 mL) per minute.

After the first bottle of 3% solution is infused and the patient shows good tolerance, subsequent infusions may be administered at a higher rate or concentration. Such increases should be made gradually allowing 15–30 minutes before each increment.

The first infusion of Panglobulin<sup>®</sup> in previously untreated agammaglobulinemic and hypogammaglobulinemic patients may lead to systemic side effects. The nature of these effects has not been fully elucidated. Some of them may be due to the release of proinflammatory cytokines by activated macrophages in immunodeficient recipients (53,54). Subsequent administration of Panglobulin<sup>®</sup> to immunodeficient patients as well as to normal individuals usually does not cause further untoward side effects.

#### Therapy of Idiopathic Thrombocytopenic Purpura (ITP)

#### Induction

0.4 g/kg of body weight on 2–5 consecutive days.

#### Acute ITP–Childhood

In acute ITP of childhood, if an initial platelet count response to the first two doses is adequate (30–50,000/μ), therapy may be discontinued after the second day of the 5 day course (17).

#### Maintenance–Chronic ITP

In adults and children, if after induction therapy the platelet count falls to less than 30,000/μl and/or the patient manifests clinically significant bleeding, 0.4 g/kg of body weight may be given as a single infusion. If an adequate response does not result, the dose can be increased to 0.8–1.0 g/kg of body weight given as a single infusion (18,55,56).


## Reconstitution

For a 3% solution using the transfer set

- Tear off the protective caps from the bottle containing the solvent and the Immune Globulin Intravenous (Human), Panglobulin<sup>®</sup>. Disinfect both rubber stoppers with alcohol.
- Remove the protective cover from one end of the transfer set and insert the needle through the rubber stopper into the bottle containing the solvent.
- Remove the cover from the other needle and plunge the inverted Panglobulin<sup>®</sup> bottle onto it, as shown in 3.
- Invert the two bottles so that the solvent flows into the Panglobulin<sup>®</sup> bottle until the required amount (see Table 2, below) has been transferred.
- Discard any unused solvent and the transfer set.

For a 6% solution using the transfer set

- Follow steps 1–3 aforementioned.
- Invert the two bottles so that the solvent flows into the Panglobulin<sup>®</sup> bottle. Use the appropriate amount (see Table 2, below) of solvent by removing the solvent bottle with transfer needle as soon as the fluid reaches the 6% mark printed on the Panglobulin<sup>®</sup> label.
- Discard any unused solvent and the transfer set.



To reconstitute Panglobulin<sup>®</sup> from the individual vial package, or when using other diluents or higher concentrations, Table 2 indicates the volume of sterile diluent required. Observing aseptic technique, this volume should be drawn into a sterile hypodermic syringe and needle. The diluent is then injected into the corresponding Panglobulin<sup>®</sup> vial size.

<b>Table 2</b>					
<b>Required Diluent Volume*</b>	<b>1 g</b>	<b>3 g</b>	<b>6 g</b>	<b>12 g</b>	
<b>Concentration</b>	<b>Vial</b>	<b>Vial</b>	<b>Vial</b>	<b>Vial</b>	<b>**</b>
3%	33.0 cc	100 cc	200 cc	200	
6%	16.5 cc	50 cc	100 cc	100	cc
9%	11.0 cc	33 cc	66 cc	132 cc	
12%	8.3 cc	25 cc	50 cc	100 cc	

\* In patients judged to be at increased risk of developing renal insufficiency, the concentration and infusion rate of Panglobulin<sup>®</sup> should be the minimum practicable.

\*\* Container not large enough to permit

this concentration.

If large doses of Panglobulin<sup>®</sup> are to be administered, several reconstituted vials of identical concentration and diluent may be pooled in an empty sterile glass or plastic i.v. infusion container using aseptic technique. Panglobulin<sup>®</sup> normally dissolves within a few minutes, though in exceptional cases it may take up to 20 minutes.

**DO NOT SHAKE! Excessive shaking will cause foaming.**

Any undissolved particles should respond to careful rotation of the bottle. Avoid foaming. Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

Filtering of Panglobulin<sup>®</sup> is acceptable but not required. Pore sizes of 15 microns or larger will be less likely to slow infusion, especially with higher Panglobulin<sup>®</sup> concentrations. Antibacterial filters (0.2 microns) may be used. When reconstitution of Panglobulin<sup>®</sup> occurs outside of sterile laminar air flow conditions, administration must begin promptly with partially used vials discarded. When reconstitution is carried out in a sterile laminar flow hood using aseptic technique, administration may begin within 24 hours provided the solution has been refrigerated during that time. Do not freeze Panglobulin<sup>®</sup> solution.

**PROCEED WITH INFUSION ONLY IF SOLUTION IS CLEAR AND AT APPROXIMATELY ROOM TEMPERATURE!**

### How supplied

Immune Globulin Intravenous (Human), Panglobulin<sup>®</sup>, is available as a white lyophilized powder in 1, 3, 6 and 12 g size vials. The only diluents which may be used to reconstitute the product are sterile (0.9%) Sodium Chloride Injection USP, 5% Dextrose, or Sterile Water. Panglobulin<sup>®</sup> is available in individual vial packages.

- 1 g Individual vial package (NDC 52769-266-61)
  - 3 g Individual vial package (NDC 52769-267-63)
  - 6 g Individual vial package (NDC 52769-268-66)
  - 12 g Individual vial package (NDC 52769-269-72)
- Please see Table 1 for Calculated Panglobulin<sup>®</sup> Osmolality (mOsm/kg).

#### Store and Dispense

Immune Globulin Intravenous (Human), Panglobulin<sup>®</sup>, should be stored at room temperature not exceeding 30 °C (86 °F). The preparation should not be used after the expiration date printed on the label.

## References

- Gardi A: Quality control in the production of an immunoglobulin for intravenous use. *Blut* 1984; 48:337–344.
- Römer J, Morgenthaler JJ, Scherz R, et al: Characterization of various immunoglobulin-preparations for intravenous application. I. Protein composition and antibody content. *Vox Sang* 1982; 42:62–73.

3. Römer J, Späth PJ, Skvaril F, et al: Characterization of various immunoglobulin preparations for intravenous application. II. Complement activation and binding to Staphylococcus protein A. *Vox Sang* 1982; 42:74–80.

- Römer J, Späth PJ: Molecular composition of immunoglobulin preparations and its relation to complement activation, in Nydegger UE (ed): Immunohemotherapy: A Guide to Immunoglobulin Prophylaxis and Therapy. London, Academic Press, 1981, pp 123–130.
- Skvaril F, Roth-Wicky B, and Barandun S: IgG subclasses in human-g-globulin preparations for intravenous use and their reactivity with Staphylococcus protein A. *Vox Sang* 1980; 38:147.
- Skvaril F: Qualitative and quantitative aspects of IgG subclasses in i.v. immunoglobulin preparations, in Nydegger UE (ed): Immunohemotherapy: A Guide to Immunoglobulin Prophylaxis and Therapy. London, Academic Press, 1981, pp 113–122.
- Skvaril F, and Barandun S: In vitro characterization of immunoglobulins for intravenous use, in Alving BM, Finlayson JS (eds): Immunoglobulins: Characteristics and Uses of Intravenous Preparations, DHHS Publication No. (FDA)-80-9005. US Government Printing Office, 1980, pp 201–206.

- Burckhardt JJ, Gardi A, Oxelius V, et al: Immunoglobulin G subclass distribution in three human intravenous immunoglobulin preparations. *Vox Sang* 1989; 57:10–14.
- Morell A, and Skvaril F: Struktur und biologische Eigenschaften von Immunoglobulinen und g-Globulin-Präparaten. II. Eigenschaften von g-Globulin-Präparaten. *Schweiz Med Wochenschr* 1980; 110:80.
- Morell A, Schürch B, Ryser D, et al: In vivo behaviour of gamma globulin preparations. *Vox Sang* 1980; 38:272.
- Imbach P, Barandun S, d’Apuzzo V, et al: High-dose intravenous gamma globulin for idiopathic thrombocytopenic purpura in childhood. *Lancet* 1981; 1:1228.
- Barandun S, Morell A, Skvaril F: Clinical experiences with immunoglobulin for intravenous use, in Alving BM, Finlayson JS (eds): Immunoglobulins: Characteristics and Uses of Intravenous Preparations. DHHS Publication No. (FDA)-80-9005. US Government Printing Office, 1980, pp 31–35.
- Schiff R, Sedlak D, Buckley R: Rapid infusion of Sandoglobulin<sup>®</sup> in patients with primary humoral immunodeficiency. *J Allergy Clin Immunol*, 88:61, 1991.
- Joller PW, Barandun S, Hitzig WH: Neue Möglichkeiten der Immunoglobulin-Ersatztherapie bei Antikörpermangel. Syndrom. *Schweiz Med Wochenschr* 1980; 110:1451.

- Barandun S, Imbach P, Morell A, et al: Clinical indications for immunoglobulin infusion, in Nydegger UE (ed): Immunohemotherapy: A Guide to Immunoglobulin Prophylaxis and Therapy. London, Academic Press, 1981, pp 275–282.
- Cunningham-Rundles C, Smithwick EM, Siegal FP, et al: Treatment of primary humoral immunodeficiency disease with intravenous (pH 4.0 treated) gamma globulin, in Nydegger UE (ed): Guide to Immunoglobulin Prophylaxis and Therapy. London, Academic Press, 1981, pp 283–290.
- Imbach P, Wagner HP, Berchtold W, et al: Intravenous immunoglobulin versus oral corticosteroids in acute immune thrombocytopenic purpura in childhood. *Lancet* 1985; 2:464.
- Fehr J, Hofmann V, Kappeler U: Transient reversal of thrombocytopenia in idiopathic thrombocytopenic purpura by high-dose intravenous gamma globulin. *N Engl J Med* 1982; 306:1254.
- Müller-Eckhardt C, Künzlen E, Thilo-Körner D, et al: High-dose intravenous immunoglobulin for posttransfusion purpura. *N Engl J Med* 1983; 308:287.
- Wenske G, Gaedicke G, Künzlen E, et al: Treatment of idiopathic thrombocytopenic purpura in pregnancy by high-dose intravenous immunoglobulin. *Blut* 1983; 46:347–353.
- Newland AC, Treleaven JG, Minchinton B, et al: High-

dose intravenous IgG in adults with autoimmune thrombocytopenia. *Lancet* 1983; 1:84–87.

- Bussel JB, Kimberly RP, Inman RD, et al: Intravenous gammaglobulin for chronic idiopathic thrombocytopenic purpura. *Blood* 1983; 62:480–486.
- Abe T, Matsuda J, Kawasugi K, et al: Clinical effect of intravenous immunoglobulin in chronic idiopathic thrombocytopenic purpura. *Blut* 1983; 47:69–75.
- Bussel JB, Schulman I, Hilgartner MW, et al: Intravenous use of gamma globulin in the treatment of chronic immune thrombocytopenic purpura as a means to defer splenectomy. *J Pediatr* 1983; 103:651–654.
- Imholz B, et al: Intravenous immunoglobulin (i.v. IgG) for previously treated acute or for chronic idiopathic thrombocytopenic purpura (ITP) in childhood: A prospective multicenter study. *Blut* 1988; 56:63–68.
- Lusher JM, and Warrior I: Use of intravenous gamma globulin in children with idiopathic thrombocytopenic purpura and other immune thrombocytopenias. *Am J Med* 1987; 83 (suppl 4A):10–16.
- Winward DB, Brophy MT: Acute renal failure after administration of intravenous immunoglobulin: Review of the literature and case report. *Pharmacotherapy* 1995; 15: 765–772.
- Cantù TG, Hoehn-Saric EW, Burgess KM, Racusen L, Scheel P: Acute renal failure associated with immunoglobulin therapy. *Am J. Kidney Dis.* 1995; 25:228–234.
- Cayco AV, Perazelly Ma, Hayslett JP: Renal insufficiency after intravenous immune globulin therapy: A Report of Two Cases and an Analysis of the Literature. *J. Amer. Soc Nephrology.* 1997; 8:1788–1793.
- Rault R, Pirano B, Johston J, Oral A: Pulmonary and renal toxicity of intravenous immunoglobulin. *Clin Nephrol:* 1991, 36:83–86.
- Michail S, Nakopoulou L, Stravrianopoulos I, Stamatiadis D, Avdikou K, Vaiopoulos G, Stathakis C: Acute renal failure associated with immunoglobulin administration. *Nephrol Dial Transplant.* 1997; 12:1497–99.
- Ashan N, Wiegand LA, Abendroth CS, Manning EC: Acute renal failure following immunoglobulin therapy. *Am J Nephrol.* 1996; 16:532–6.
- Hammarstrom L, and Smith CI: Placental transfer of intravenous immunoglobulin. *Lancet* 1986; 1:681.
- Sidiropoulos D, et al: Transplacental passage of intravenous immunoglobulin in the last trimester of pregnancy. *J Pediatr* 1986; 109:505–508.
- Wenske G, et al: Idiopathic thrombocytopenic purpura in pregnancy and neonatal period. *Blut* 1984; 48:377–382.
- Fabris P, et al: Successful treatment of a steroid-resistant form of idiopathic thrombocytopenic purpura in pregnancy with high doses of intravenous immunoglobulins. *Acta Haemat* 1987; 77:107–110.
- Coller BS, et al: Management of severe ITP during pregnancy with intravenous immunoglobulin (IVIgG). *Clin Res* 1985; 33:545A.
- Tchernia G, et al: Management of immune thrombocytopenia in pregnancy: Response to infusions of immunoglobulins. *Am J Obstet Gynecol* 1984; 148:225–226.
- Newland AC, et al: Intravenous IgG for autoimmune thrombocytopenia in pregnancy. *N Engl J Med* 1984; 310:261–262.
- Morgenstern GR, et al: Autoimmune thrombocytopenia in pregnancy: New approach to management. *Br Med J* 1983; 287:584.
- Ciccimarra F, et al: Treatment of neonatal passive immune thrombocytopenia. *J Pediat* 1984; 105:677–678.
- Rose VL, and Gordon LI: Idiopathic thrombocytopenic purpura in pregnancy. Successful management with immunoglobulin infusion. *JAMA* 1985; 254: 2626–2628.
- Gounder MP, et al: Intravenous gammaglobulin therapy in the management of a patient with idiopathic thrombocytopenic purpura and a warm autoimmune erythrocyte panagglutinin during pregnancy. *Obstet Gynecol* 1986; 67:741–746.
- Siber GR, Werner BG, Halsey NA, et al: Interference of immune globulin with measles and rubella immunisation. *J. Pediatr.* 1993; 122:204–211.

45. American Academy of Pediatrics, Committee on Infectious Diseases: Recommended timing of routine measles immunization for children who have recently received immune globulin preparations. *Pediatrics* 1994; 93:682–685.

46. Centers of Disease Control and Prevention Measles, mumps, and rubella-vaccine use and strategies for elimination of measles, rubella, and congenital rubella syndrome and control of mumps: recommendations of the advisory committee on immunization practices (ACIP). *MMWR, Morbidity and Mortality Weekly Report.* May 22, 1998; vol 47/No. RR-8, 1–57.

47. Phillips AO: Renal failure and intravenous immunoglobulin [letter; comment]. *Clin Nephrol* 1992; 36: 83–86.

48. Cunningham-Rundles C, Day NK, Wahn V, et al: Reactions to intravenous gamma globulin infusions and immune complex formation, in Nydegger UE (ed): Immunohemotherapy: A Guide to Immunoglobulin Prophylaxis and Therapy. London, Academic Press, 1981, pp 447–449.

49. Brox AG, Cournoyer D, et al: Hemolytic anemia following intravenous gamma globulin administration. *Am J. Med.* 1987; 82:633–635.

50. Kim HC, Park CL, Cowan JH, et al: Massive intravascular hemolysis associated with intravenous immunoglobulin in bone marrow transplant recipients. *Am J. Ped. Hematol/Oncol* 1988; 10:67–74.

51. Nicholls MD, Cummins JC, et al: Haemolysis induced by intravenously-administered immunoglobulin. *Med J of Australia* 1989; 150:404–406.

52. Copelan EA et al: Hemolysis following intravenous immune globulin therapy. *Transfusion* 1986; 26:410–412.

53. Aukrust P, Froland SS, N-B Liabakk, F. Müller, et al: Release of cytokines, soluble cytokine receptors, and interleukin-1 receptor antagonist after intravenous immunoglobulin administration in vivo. *Blood* 1994; 84:2136–2143.

54. Bagdasarian A, Tonetta S, Harel W, Mamidi R. Uemura Y: IVIG adverse reactions: potential role of cytokines and vasoactive substances. *Vox Sang* 1998; 74:74–82.

55. Bussel JB, Pham LC, Hilgartner MW, et al: Long-term maintenance of adults with ITP using intravenous gamma globulin. Abstract, America Society of Hematology, New Orleans, December, 1985.

56. Imbach PA, Kühne T, Holländer G: Immunologic aspects in the pathogenesis and treatment of immune thrombocytopenic purpura in children. *Current opinion in Pediatrics* 1997; 9:35–40.

57. Anderson W, Bethea W: Renal lesions following administration of hypertonic solutions of sucrose. *JAMA.* 1940; 114:1983–1987.

58. Lindberg H, Wald A: Renal lesions following the administration of hypertonic solutions: *Arch Intern Med.* 1939; 63:907–918.

59. Rigdon RH, Cardwell ES: Renal lesions following the intravenous injection of hypertonic solution of sucrose: A clinical and experimental study. *Arch Intern Med.* 1942; 69:670–690.

**Manufactured by:**  
**ZLB Bioplasma AG**  
**Wankdorfstrasse 10, 3000 Berne 22**  
**Switzerland**  
**US License No. 1598**

**Distributed by:**  
**AMERICAN RED CROSS**  
**BLOOD SERVICES**  
**WASHINGTON, D.C. 20006, USA**

**September 00**